

	MEDIC	CAL HISTORY		
Name	Ad	dress		
City	State	Zip	_Email	
HomePhone	W	ork/Cell Phone		
Primary Physician's Name		Ph	one #	
Primary Physician's NameB/PTP	RDOB	Age	Ht	Wt
Please list all medications you	are currently takir	ng:		
Allergies:	Are you o	n Antibiotics at this	time?	
Circle any of the following illne	sses you have or	have ever had in th	e past:	
Myesthenia Gravis Hepatitis Numbness Muscle Weakne Parkinson's Disease Neuro List and/or Explain Other Medicabove:	Eye Disease ess Multiple S logical Disorders cal Conditions not	Autoimmune E clerosis Amyotro Lambert-Eato listed	Disease Vi ophic Lateral S n Syndrome	
PreviousHospitalizations/Opera				
WOMEN: Are you Pregnant, Tryi	ing to get Pregnai	nt, or Lactating (nur		
Have you had Plastic Surgery o	or other surgery to	your face/neck are	eas when?	
Had Botox® injections before?	Last treat	ment?V	Vhat Areas?	
ExplainWe		-	eatments?	
Have you ever had eyelid/eye	brow droop after	Botox®?		
Do you show a lot of upper eye	e lid when eyes ar	e open?		
Do your eyelids feel extra heav		get enough sleep?	?	
Do your eyelids droop without s	•			
Areas of special concern to pa	itient?			
I understand the information on the provision of treatment. I u report it to the office as soon questionnaire. I acknowledge member responsible for any en	nderstand that if as possible. I h that all answers h	any changes occu ave read and und ave been recorded	r in my medio lerstand the o d truthfully and	cal history/health I wi above medical histor d will not hold any sta
Patient Signature			Date	



CONSENT TO BOTOX® BOTULINUM TOXIN "A" TREATMENT

Botox® a neurotoxin produced by the bacterium Clostridium A. Botox® can relax the muscles on areas of the face and neck which cause wrinkles associated with facial expressions. Treatment with Botox can cause your facial expression lines or wrinkles to essentially disappear. Areas most frequently treated are: a) glabellar area of frown lines, located between the eyes; b) crow's feet (lateral areas of the eyes); and c) forehead wrinkles. Botox is diluted to a very controlled solution and when injected into the muscles with a very thin needle, it is almost painless. Clients may feel a slight burning sensation while the solution is being injected. The procedure takes about 15-20 minutes and the results last 3-6 months. With repeated treatments, the results may tend to last longer.

RISKS AND COMPLICATIONS

It has been explained to me that there are certain inherent and potential risks and side effects in any invasive procedure and in this specific instance such risks include but are not limited to: 1.Post treatment discomfort, swelling, redness, and bruising, 2.Double Vision 3. Rarely weakened tear duct 5. Post treatment bacterial, and/or fungal infection requiring further treatment 6.Allergic reaction 7. Minor temporary droop of eyelid(s) in approximately 2% of injections, this usually lasts 2-3 weeks 8. Occasional numbness of the forehead lasting up to 2-3 weeks, 9.Transient headache, and 10. Flu-like symptoms may occur.

PHOTOGRAPHS

I authorize the taking of clinical photographs and their use for scientific purposes both in publications and presentations. I understand my identity will be protected.

PREGNANCY, ALLERGIES & NEUROLOGIC DISEASE

I am not aware that I am pregnant and I am not trying to get pregnant, I am not Lactating (nursing), have any significant Neurologic disease including but not limited to Myasthenis Gravis, Multiple Sclerosis, Lambert-Eaton Syndrome, Amyotrophic Lateral Sclerosis (ALS), Parkinson's or that I have any allergies to the toxin ingredients, or to human albumin.

PAYMENT

I understand that this is an "elective" cosmetic procedure and that payment is my responsibility.

RESULTS

I am aware that when small amounts of purified botulinum ("BOTOX®") are injected into a muscle it causes weakness or paralysis of that muscle. This appears in 2 – 10 days and usually lasts 3-6 months but can be shorter or longer. In a very small number of individuals, the injection does not work as satisfactorily or for as long as usual and their some individuals who do not respond at all. I understand that I will not be able to "frown" while the injection is effective but that this will reverse after a period of months at which time re-treatment is appropriate. I understand that I must stay in the erect posture and that I must not manipulate the area (s) of the injections for the 2 hours post-injection period.

I understand this an elective procedure and I hereby voluntarily consent to treatment with Botox® injection for the condition known as: Facial Dynamic Wrinkles. The procedure has been fully explained to me. I have read the above and understand it. My questions have been answered satisfactorily. I accept the risks and complications of the procedure and I understand that no guarantees are implied as to the outcome of the procedure. I also certify that if I have any changes in my medical history I will notify the office immediately.

Patient Name (Print)	Patient Signature	Date	
Witness Name (Print)	Witness Signature	Date	



PRE - TREATMENT INSTRUCTIONS

In an ideal situation it is prudent to follow some simple guidelines before treatment that can make all the difference between a fair result or great result, by reducing some possible side effects associated with the injections. We realize this is not always possible; however, minimizing these risks is always desirable.

- Avoid Alcoholic beverages at least 24 hours prior to treatment (Alcohol may thin the blood increasing risk of bruising.
- Avoid Anti-inflammatory / Blood Thinning medications ideally, for a
 period of two (2) weeks before treatment. Medications and
 supplements such as Aspirin, Vitamin E, Gingo Biloba, St. John's
 Wort, Ibuprofen, Motrin, Advil, Aleve, Vioxx, and other NSAIDS are all
 blood thinning and can increase the risk of bruising/swelling after
 injections.
- Schedule Botox® appointment at least 2 weeks prior to a special event which may be occurring, i.e., wedding, vacation, etc. etc. It is not desirable to have a very special event occurring and be bruised from an injection which could have been avoided.



BOTOX® POST - TREATMENT INSTRUCTIONS

The guidelines to follow post treatment have been followed for years, and are still employed today to prevent the possible side effect of ptosis. These measures should minimize the possibility of ptosis almost 98%.

- No straining, heavy lifting, vigorous exercise for 3-4 hours following treatment. It is now known that it takes the toxin approximately 2 hours to bind itself to the nerve to start its work, and because we do not want to increase circulation to that area to wash away the Botox® from where it was injected. This waiting period continues to be recommended by most practitioners.
- Avoid Manipulation of area for 3-4 hours following treatment. (For the same reasons listed above.) This includes not doing a facial, peel, or micro-dermabrasion after treatment with Botox®. A facial, peel, or micro-dermabrasion can be done in same appointment only if they are done before the Botox®.
- Facial Exercises in the injected areas is recommended for 1-hour following treatment, to stimulate the binding of the toxin only to this localized area.
- Do not lie down or bend over for 3-4 hours following treatment. (This instruction has been employed for years by some practitioners, although, we have not been able to find out the main reason for this since many practitioners inject while the patient is in a lying position. Many practitioners do not adhere to this anymore.
- It can take 2-10 days to take full effect. It is recommended that the
 patient contact office no later than 2 weeks after treatment if
 desired effect was not achieved and no sooner to give toxin time to
 work.

Makeup may be applied before leaving the office. Some practitioners recommend avoiding Retin-A, Glycolic acid, Vitamin C, and Kinerase for 24 hrs to the treated areas.